

**CANADIAN BOARD OF EXAMINERS FOR PROFESSIONAL SURVEYORS
ATLANTIC PROVINCES BOARD OF EXAMINERS FOR LAND SURVEYORS**

SCHEDULE I / ITEM 2

October 2008

LEAST SQUARES ESTIMATION AND DATA ANALYSIS

Note: This examination consists of 8 questions on 3 pages.



Marks

Q. No

Time: 3 hours

Value Earned

1	Define or explain the following terms:	2	
	a) Precision	2	
	b) Accuracy	2	
	c) Standard deviation	2	
	d) Root mean square error	2	
	e) Correlation coefficient	2	
	f) Redundancy of a linear system	2	
	g) Type I and type II errors in statistical testing	2	
2	Given the cofactor matrix Q of the horizontal coordinate (x, y) of a survey station and the unit variance $\hat{\sigma}_0^2 = 2 \text{ cm}^2$, calculate the semi-major, semi-minor axis and the orientation of the standard error ellipse associated with this station.	10	
	$Q = \begin{bmatrix} 5.32 & 6.02 \\ 6.02 & 8.38 \end{bmatrix}$		
3	Given the following mathematical model		
	$f(\lambda, x) = 0 \quad C_\lambda \quad C_x$		
	where f is the vector of mathematical models, x is the vector of unknown parameters and C_x is its variance matrix, λ is the vector of observations and C_λ is its variance matrix.		
	a) Linearize the mathematical model	2	
	b) Formulate the variation function	3	
	c) Derive the least squares normal equation	4	
	d) Derive the least squares solution of the unknown parameters.	6	

4	<p>Given a leveling network below where A and B are known points, h_1 and h_2 are two height difference measurements with standard deviation of σ_1 and σ_2, respectively and $\sigma_1 = 2 \sigma_2$. Determine the value of σ_1 and σ_2 so that the standard deviation of the height solution at P using a least squares adjustment is equal to 2 mm.</p> 	10													
5	<p>Given the variance-covariance matrix of the measurement vector $\lambda = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix}$:</p> $C_\lambda = \begin{bmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{bmatrix}$ <p>and two functions of λ: $x = \lambda_1 + \lambda_2$ and $y = 3\lambda_1$, determine $C_{xy}, C_{x\lambda}, C_{y\lambda}$</p>	10													
6	<p>Given the angle measurements at a station along with their standard deviations:</p> <table border="1" data-bbox="425 1138 1198 1285"> <thead> <tr> <th>Angle</th> <th>Measurement</th> <th>Standard Deviation</th> </tr> </thead> <tbody> <tr> <td>α</td> <td>$134^\circ 38' 56''$</td> <td>$6.7''$</td> </tr> <tr> <td>β</td> <td>$83^\circ 17' 35''$</td> <td>$9.9''$</td> </tr> <tr> <td>γ</td> <td>$142^\circ 03' 14''$</td> <td>$4.3''$</td> </tr> </tbody> </table>  <p>Perform least squares adjustment to the problem using</p> <ol style="list-style-type: none"> Conditional equations (conditional adjustment) Observation equations (parametric adjustment) 	Angle	Measurement	Standard Deviation	α	$134^\circ 38' 56''$	$6.7''$	β	$83^\circ 17' 35''$	$9.9''$	γ	$142^\circ 03' 14''$	$4.3''$	25	
Angle	Measurement	Standard Deviation													
α	$134^\circ 38' 56''$	$6.7''$													
β	$83^\circ 17' 35''$	$9.9''$													
γ	$142^\circ 03' 14''$	$4.3''$													

7	<p>Given the sample unit variance obtained from the adjustment of a geodetic network $\hat{\sigma}_0^2 = 0.55 \text{ cm}^2$ with a degree of freedom $\nu = 3$ and the a-priori standard deviation $\sigma_0 = 0.44 \text{ cm}$, conduct a statistic test to decide if the adjustment result is acceptable with a significance level of $\alpha = 5\%$.</p> <p>The critical values that might be required in the testing are provided in the following table:</p> <table border="1" data-bbox="461 537 1174 642"> <tr> <td>α</td> <td>0.001</td> <td>0.01</td> <td>0.025</td> <td>0.05</td> <td>0.10</td> </tr> <tr> <td>$\chi_{\alpha, \nu=3}^2$</td> <td>16.26</td> <td>11.34</td> <td>9.35</td> <td>7.82</td> <td>6.25</td> </tr> </table> <p>where $\chi_{\alpha, \nu=3}^2$ is determined by the equation $\alpha = \int_{\chi_{\alpha, \nu=3}^2}^{\infty} \chi^2(x) dx$.</p>	α	0.001	0.01	0.025	0.05	0.10	$\chi_{\alpha, \nu=3}^2$	16.26	11.34	9.35	7.82	6.25	5					
α	0.001	0.01	0.025	0.05	0.10														
$\chi_{\alpha, \nu=3}^2$	16.26	11.34	9.35	7.82	6.25														
8	<p>The following residual vector \hat{r} and estimated cofactor matrix $Q_{\hat{r}}$ were computed from a least squares adjustment using independent observations with a standard deviation of $\sigma_0 = 1.5 \text{ mm}$:</p> $\hat{r} = [4 \quad 2 \quad -3 \quad 10] \quad (\text{mm})$ $Q_{\hat{r}} = \begin{bmatrix} 15 & 1 & 3 & -2 \\ 1 & 7 & -1 & 3 \\ 3 & -1 & 4 & -1 \\ -2 & 3 & -1 & 2 \end{bmatrix} \quad (\text{mm}^2)$ <p>Given that a global test has been rejected with a significance level of $\alpha = 0.04$, conduct further tests to identify which observation(s) may contain an outlier. The critical values that might be required in the testing are provided in the following table:</p> <table border="1" data-bbox="371 1369 1263 1459"> <tr> <td>α</td> <td>0.001</td> <td>0.002</td> <td>0.003</td> <td>0.004</td> <td>0.005</td> <td>0.01</td> <td>0.05</td> </tr> <tr> <td>K_{α}</td> <td>3.09</td> <td>2.88</td> <td>2.75</td> <td>2.65</td> <td>2.58</td> <td>2.33</td> <td>1.64</td> </tr> </table> <p>where K_{α} is determined by the equation $\alpha = \int_{K_{\alpha}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$.</p>	α	0.001	0.002	0.003	0.004	0.005	0.01	0.05	K_{α}	3.09	2.88	2.75	2.65	2.58	2.33	1.64	10	
α	0.001	0.002	0.003	0.004	0.005	0.01	0.05												
K_{α}	3.09	2.88	2.75	2.65	2.58	2.33	1.64												
Total Marks:			100																